arXiv:1103.5547v1 [math.FA] 29 Mar 2011

A NOTE ON HOMOGENEOUS FUNCTIONS AND PARALLELOGRAM LAW

WENHAN WANG AND WEN WANG

Abstract. In this article, we investigate a new characterization of the parallelogram law in a normed linear space. We give equivalent conditions to the parallelogram law, in terms of the homogeneous property of a continuous positive definite function on the normed space.

Let $K = \mathbf{R}$ or $\mathbf{C}$ be the base field, and suppose X be a normed K-linear space, with norm $\|\cdot\|$. A continuous function $p : X \to \mathbf{R}$ is called *positive definite* if $p(x) \geq 0$ for all $x \in X$, and $p(x) = 0$ if and only if $x = 0 \in X$. Moreover, $p : X \to \mathbf{R}$ is called homogeneous, if for any $\lambda \in K$, $p(\lambda x) = |\lambda|\, p(x)$. We first note the following lemma on the comparison of two positive definite homogeneous continuous functions on X.

Lemma 1. *Let $p : X \to \mathbf{R}$ be a continuous homogeneous function on X. If p is bounded, then $p \equiv 0$ on X.*

Proof. Since p is bounded, we may assume that the $|p(x)| \leq M$ for all $x \in X$. It then follows that for any $x \in X$ and any $\varepsilon > 0$,

$$|p(x)| = \varepsilon \cdot \left| p\left(\frac{x}{\varepsilon}\right)\right| \leq \varepsilon \cdot M.$$

Since $\varepsilon > 0$ is arbitrary, it is necessary that $|p(x)| = 0$. Hence $p(x) = 0$. □

We then obtain immediately the following corollary on the comparison of two homogeneous functions $p_1(x)$ and $p_2(x)$.

Corollary 2. *Let p_1 and p_2 be two homogeneous functions on X, such that $p_1 - p_2$ is bounded on X. Then $p_1 = p_2$.*

Proof. Apply the above Lemma to $p = p_1 - p_2$. □

We then introduce and prove the main Theorem on characterization of the parallelogram law in terms of homogeneous functions. For any $y \in X$ with $\|y\| = 1$, we define a function

$$p_y(x) := \sqrt{\frac{1}{2}\left(\|x+y\|^2 + \|x-y\|^2\right) - 1}. \tag{1}$$

Note that $p_y : X \to \mathbf{R}$ is well-defined. To see this, note that

$$\|x+y\|^2 + \|x-y\|^2 \geq \frac{1}{2}\left(\|x+y\| + \|x-y\|\right)^2 \geq \frac{1}{2}\|2y\|^2 = 2,$$

hence $\frac{1}{2}\left(\|x+y\|^2+\|x-y\|^2\right)-1\geq 0$, and p_y is well-defined from X to $\mathbf{R}$. Moreover, following a closer look at the above arguments, we have

Proposition 3. *For any $y\in X$, p_y is a continuous positive definite function.*

Before we introduce the main Theorem, we shall compare $p_y(x)$ with the norm $\|x\|$.

Lemma 4. *For any $x\in X$, with $\|x\|\geq 1$, we have $\|x\|-1\leq p_y(x)\leq \|x\|+1$.*

Proof. To show that $p_y(x)\geq \|x\|-1$, note that

$$\begin{array}{rcl} p_y(x) & = & \sqrt{\frac{1}{2}\left(\|x+y\|^2+\|x-y\|^2\right)-1} \\ & \geq & \sqrt{\frac{1}{4}\left(\|x+y\|+\|x-y\|\right)^2-1} \\ & \geq & \sqrt{\frac{1}{4}\|2x\|^2-1} \\ & \geq & \sqrt{\|x\|^2-1} \\ & \geq & \sqrt{\|x\|^2-2\|x\|+1} \\ & = & \|x\|-1. \end{array}$$

To prove the other inequality, note that

$$\begin{array}{rcl} p_y(x) & = & \sqrt{\frac{1}{2}\left(\|x+y\|^2+\|x-y\|^2\right)-1} \\ & \leq & \sqrt{\|x\|^2+2\|x\|\|y\|+\|y\|^2-1} \\ & = & \sqrt{\|x\|^2+2\|x\|} \\ & < & \sqrt{\|x\|^2+2\|x\|+1} \\ & = & \|x\|+1. \end{array}$$

□

Now we may give the main Theorem, on the characterization of parallelogram rule in terms of $p_y(x)$, for all $y\in X$.

Theorem 5. *Let X be a normed linear space, and for each $y\in X$, $p_y(x)$ is defined as above. Then the following statements are equivalent.*

(i) *For all $y\in X$, p_y is homogeneous.*
(ii) *For all $y\in X$, $p_y(x)=\|x\|$.*
(iii) *$\|\cdot\|$ is induced from an inner product on X.*

Proof. (i)⇒(ii). Note that both p_y and the norm $\|\cdot\|$ are homogeneous functions on X. The above lemma shows that $p_y(x) - \|x\|$is bounded on X. Hence $p_y(x) = \|x\|$ for all $x \in X$.
(ii)⇒(iii). Note that $p_y(x) = \|x\|$ gives that for any $y \in X$ with $\|y\| = 1$, $2\|x\|^2 + 2\|y\|^2 = \|x+y\|^2 + \|x-y\|^2$. In general, if $y \neq 0$, then this also implies the parallelogram law, by taking $\tilde{y} = \frac{y}{\|y\|}$.
(iii)⇒(i). If $\|\cdot\|$ is induced from an inner product $\langle \cdot, \cdot \rangle : X \times X \to \mathbf{C}$, then we may prove directly that $p_y(x) = \langle x, x \rangle$, which is thus homogeneous. □

References

1. Barbara D. MacCleur. *Elementary Functional Analysis.* Springer, New York, London, 2008.

E-mail address: `wangwh@math.washington.edu`

E-mail address: `wwang@math.wsu.edu`